

ACCEPTED MANUSCRIPT • OPEN ACCESS

## Recent peat and carbon accumulation following the Little Ice Age in northwestern Québec, Canada

To cite this article before publication: Sanna Riikka Piilo *et al* 2019 *Environ. Res. Lett.* in press <https://doi.org/10.1088/1748-9326/ab11ec>

### Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2018 The Author(s). Published by IOP Publishing Ltd.

As the Version of Record of this article is going to be / has been published on a gold open access basis under a CC BY 3.0 licence, this Accepted Manuscript is available for reuse under a CC BY 3.0 licence immediately.

Everyone is permitted to use all or part of the original content in this article, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions may be required. All third party content is fully copyright protected and is not published on a gold open access basis under a CC BY licence, unless that is specifically stated in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

1

2

31Recent peat and carbon accumulation following the Little Ice Age in northwestern Québec, Canada

4

5

62Sanna R. Piilo<sup>\*1,2</sup>, Hui Zhang<sup>1,2</sup>, Michelle Garneau<sup>3</sup>, Angela Gallego-Sala<sup>4</sup>, Matthew J. Amesbury<sup>1,4</sup>, Minna M.

7

8Väiliranta<sup>1,2</sup>

9

104Abstract

11

12

135Peatland ecosystems are important carbon sinks, but also release carbon back to the atmosphere as carbon dioxide and

14

156methane. Peatlands therefore play an essential role in the global carbon cycle. However, the response of high-latitude

16

177peatlands to ongoing climate change is still not fully understood. In this study, we used plant macrofossils and peat

18

198property analyses as proxies to document changes in vegetation and peat and carbon accumulation after the Little Ice

20

219Age. Results from 12 peat monoliths collected in high-boreal and low-subarctic regions in northwestern Québec,

22

2310Canada, suggest high carbon accumulation rates for the recent past (post AD 1970s). Successional changes in plant

24

2511assemblages were asynchronous within the cores in the southernmost region, but more consistent in the northern region.

26

2712Average apparent recent carbon accumulation rates varied between 50.7 and 149.1 g C m<sup>-2</sup> yr<sup>-1</sup> with the northernmost

28

2913study region showing higher values. The variation in vegetation records and peat properties found within samples taken

30

3114from the same sites and amongst cores taken from different regions highlights the need to investigate multiple records

32

3315from each peatland, but also from different peatlands within one region.

34

3516<sup>1</sup>ECRU, Ecosystems and Environment Research Programme, Department of Environmental Sciences, University of

36

3717Helsinki, P.O. Box 65, 00014, Finland

38

39

4018<sup>2</sup> Helsinki Institute of Sustainability Science (HELSUS), Finland

41

42

4319<sup>3</sup> Department of Geography, Université du Québec à Montréal, Canada

44

4520<sup>4</sup> Geography, College of Life and Environmental Sciences, University of Exeter, UK

46

47

4821\*Corresponding author: email: [sanna.piilo@helsinki.fi](mailto:sanna.piilo@helsinki.fi)

49

5022**Keywords:** permafrost peatlands, climate warming, vegetation dynamics, carbon accumulation, plant macrofossil

51

5223analysis

53

5424

55

56

5725

58

59

6026

## 27 Introduction

28 Future changes in climate can be expected to influence peatland vegetation and related carbon dynamics, especially in  
29 the high-latitudes where warming is most pronounced, almost twice the global average (IPCC, 2013). Permafrost  
30 peatlands are especially sensitive to climate change, namely changes in temperatures and moisture balance  
31 (precipitation minus evaporation) which determine vegetation and carbon cycling processes (Ovenden, 1990; Carroll  
32 and Crill, 1997; Davidson and Janssens, 2006; Swindles *et al.*, 2015; Galka *et al.*, 2017, 2018). Despite covering only ca.  
33 5% of Earth's landmass (Yu *et al.*, 2010), peatlands form a major terrestrial carbon sink (Gorham, 1991; Yu *et al.*, 2009,  
34 2010) and are connected to the global carbon cycle and its associated carbon-climate feedbacks (Frolking and Roulet,  
35 2007; McGuire *et al.*, 2009; Yu, 2011). However, there are still large uncertainties about future climate-induced  
36 changes in peatland dynamics driven by changes in vegetation assemblages and moisture conditions (McGuire *et al.*,  
37 2018).

38 Due to climate warming, net primary productivity (NPP) will likely increase because of longer growing seasons  
39 (Gallego-Sala *et al.*, 2018). This could accelerate peat accumulation, an effect that may be partly, or even wholly,  
40 mitigated by an increase in peat decomposition (Ise *et al.*, 2008; Dorrepaal *et al.*, 2009; Crowther *et al.*, 2016). Yu *et al.*  
41 (2009) and Charman *et al.* (2013) have shown that during past warm periods increases in NPP exceeded increases in  
42 peat decomposition, leading to higher rates of peat accumulation. The overall predicted 30% increase in precipitation  
43 (Collins *et al.*, 2013) may also enhance peat accumulation if not reversed by increase in evapotranspiration (Yu *et al.*,  
44 2009).

45 In the James Bay and Hudson Bay Lowlands, peatlands cover vast areas and have stored ca. 30 Pg carbon since their  
46 initiation during the mid-Holocene (Packalen *et al.*, 2014). Many previous studies have investigated Holocene carbon  
47 dynamics in eastern Canada (e.g. Charman *et al.* 1994; van Bellen *et al.*, 2011a, 2011b; Garneau *et al.* 2014; Magnan  
48 and Garneau 2014a, 2014b; Packalen and Finkelstein 2014; Charman *et al.*, 2015) showing high regional variability in  
49 both apparent rates of carbon accumulation and in the timing of shifts in carbon accumulation throughout the Holocene.  
50 Studies focusing on more recent carbon dynamics and response to ongoing rapid climate change in northwestern  
51 Québec are sparse (see, however, Ali *et al.*, 2008; Beaulieu-Audy *et al.*, 2009; Loisel and Garneau, 2010; Lamarre *et al.*  
52 2012).

53 To address the knowledge gap on the developmental and carbon dynamics of high-latitude peatlands after the Little Ice  
54 Age (LIA) (anno Domini (AD) 1450-1850 (Naulier *et al.*, 2015; Wilson *et al.*, 2016)), we examined several peat  
55 sections collected from two ecoregions (high-boreal and low-subarctic), in northwestern Québec. Climate started to

warm after the LIA and since the 1970s human-induced warming has been more prominent (PAGES 2k Consortium, 2013; Abram *et al.* 2016). Increased snowfall since the 1950s and warm temperatures in the 1990s triggered rapid permafrost melting on the eastern coast of Hudson Bay (Payette *et al.*, 2004). Hence, permafrost is predicted to disappear from subarctic Québec in the coming decades (Payette *et al.*, 2004; Jean and Payette, 2014), likely altering the vegetation and thus peatland carbon dynamics of the region (Swindles *et al.*, 2015).


This study investigates the response of northern peatland vegetation and carbon accumulation dynamics to recent warming in northwestern Québec. Plant macrofossil analysis was used as a primary proxy to reconstruct past habitat changes (cf. Väliranta *et al.*, 2007; 2017).  $^{14}\text{C}$  and  $^{210}\text{Pb}$  chronologies in combination with peat property analyses enabled the reconstruction of peat and carbon accumulation. Replicate records from each peatland and from proximal locations inside the study regions allowed us to estimate if detected changes were climate-induced or driven by internal dynamics (Swindles *et al.*, 2012; Mathijssen *et al.*, 2016, 2017; Magnan *et al.*, 2018). Specifically we evaluated: 1) if plant assemblages are changing synchronously in response to rapid recent warming, 2) if warmer climate has increased peat plant net primary productivity and thus carbon accumulation rates, especially in the southern range of the study region, and 3) if multiple cores provide better assessment of within and between site variability.

## Methods

### *Study sites*

We selected two regions in northwestern Québec: one within the sporadic and the other within the discontinuous permafrost zone (Allard and K.-Seguin, 1987; Thibault and Payette, 2009). Kuujjuarapik (K1P and K2) represents the subarctic, forested tundra ecoregion at the southernmost limit of the discontinuous permafrost zone (Fig. 1). Palsa mounds are characteristic at K1P peatland, whereas K2 is a small fen with peat thickness of 1–2 metres. Radisson (LG2 and Rad) represents the northernmost boreal ecoregion (Fig. 1). These two peatlands, in region of sporadic permafrost, are ombrotrophic with peat thickness of 3–4 meters (Thibault and Payette, 2009).

In the two study regions, mean annual temperatures are below zero °C (Table 1). Mean annual temperatures have increased in both regions (Table 1). In Kuujjuarapik mean summer temperature has increased 1.0°C since 1961, while mean winter temperature has increased 0.5°C since 1971 (Environment Canada, 2018). In Radisson, since 1971, the biggest seasonal increase of 0.5°C is in mean autumn temperatures. In Kuujjuarapik, the increase in growing degree-days (GDD0) (between 1971–2000 and 1981–2010) is ca. 6% and in Radisson ca. 4% (Table 1). In both regions, autumn rainfall especially has increased. 35–40% of the precipitation falls as snow.



**Figure 1.** Study site locations marked with dots (Kuujjuarapik and Radisson). Pictures from the coring sites, (a): K1P, (b): K2, (c): Rad, and (d): LG2. Basemap: Esri, HERE, Garmin © OpenStreetMap.

In the James Bay lowlands, peatlands cover ca. 20-30% of the land area (Arseneault and Sirois, 2004). In Radisson, peat started to accumulate following post glacial land uplift after 8000 calibrated (cal.) yr. before present (BP) (Dyke and Prest, 1987; Beaulieu-Audy *et al.* 2009). Radisson study sites are characterised by hummock-hollow microforms with e.g. *Sphagnum lindbergii* and *Warnstorfia exannulata* group abundantly present in low lawns and hollows and *S. fuscum* on hummocks while *Picea mariana* is sparsely present (Table 1). Along the Hudson Bay lowlands, at the K1P, peat started to accumulate over marine sediments around 5950-5100 cal. BP and palsa formation is connected to the LIA cooling (Arlen-Pouliot and Bhiry, 2005; Lamarre *et al.* 2012). Ombrotrophic habitat conditions prevail in the central palsa mound part and hummock vegetation in K1P and K2 is largely similar to Radisson peatlands (Table 1) but supplemented by e.g. *Polytrichum strictum*, and *Sphagnum russowii*. Sedges dominate the fen habitats together with *Myrica gale*, *Menyanthes trifoliata*, *Eriophorum russeolum*, *S. papillosum*, *S. riparium*, and brown mosses such as *Scorpidium scorpioides*. *Larix laricina* is occasionally present.

**Table 1.** Study site information. Meteorological data for 30-year (1981-2010) measuring period: mean annual temperature (MAT), mean annual precipitation (MAP), and growing degree-days above zero (GDD0), measured from stations; La Grande Riviere A in Radisson and Kuujjuarapik station A (Environment Canada, 2018). Increases in MAT, MAP, and GDD0 since 1961 and/or 1971 until present. Meters above sea level (MASL). Dominant vegetation around the coring points listed.

Location	MAT (°C)	MAP (mm)	GDD0	Core name	Latitude (N)	Longitude (W)	MASL (m)	Dominant vegetation
<b>Kuujjuarapik</b>				K1P1	55°13'35.8"	77°41'41.2"	100	<i>Sphagnum fuscum</i> , <i>S.</i>
1981-2010:	-4.0	661	1384	K1P2	55°13'34.9"	77°41'47.8"		<i>capillifolium</i> , <i>Cladonia spp.</i> ,
Increase since;				K1P3	55°13'34.0"	77°41'54.7"		<i>Chamaedaphne calyculata</i> ,
1961until present:	0.5	46		K2.1	55°13'38.6"	77°42'18.4"	115	<i>Kalmia polifolia</i> ,
1971until present:	0.4	12	74	K2.2	55°13'38.2"	77°42'19.5"		<i>Rhododendron groenlandicum</i>
				K2.3	55°13'38.2"	77°42'20.6"		<i>Andromeda polifolia</i> , <i>Vaccinium</i>
<b>Radisson</b>				LG2.1	53°39'09.4"	77°44'00.2"	175	<i>uliginosum</i> , <i>V. oxycoccus</i> ,
1981-2010:	-2.9	697	1684	LG2.2	53°39'08.6"	77°43'56.2"		<i>Trichophorum cespitosum</i> ,
Increase since;				LG2.3	53°39'04.1"	77°43'50.3"		<i>Carex spp.</i> , <i>Rubus</i>
1971until present:	0.2	13	62	Rad1	53°39'38.6"	77°44'51.9"	170	<i>chamaemorus</i> , <i>Drosera</i>
				Rad2	53°39'42.0"	77°44'52.9"		<i>rotundifolia</i>
				Rad3	53°39'45.1"	77°44'52.7"		

### Sampling

In early July 2017, twelve peat monoliths: three from each four peatlands, K1P, K2, LG2, and Rad, were collected (Table 1, Fig. 1). The core-lengths varied between 32 and 39 cm (Table 2). Peat sections (10 cm in diameter) were sampled by hand sawing the layer overlying seasonal frost or permafrost from intermediate lawn microforms (water tables from 13 to 23 cm) inhabited mainly by *Sphagnum fuscum*. Such microhabitats are considered the most sensitive to reflect climate-induced changes in hydrology and consequent plant community shifts (e.g. Välranta *et al.*, 2007; Frolking *et al.*, 2014). The cores were collected from relatively central parts of the peatlands, some tens of meters apart from each other. Peat monoliths were wrapped in plastic film and aluminium foil, transported and frozen at the University of Helsinki, Finland. For the analyses, monoliths were cut into 1-cm slices. To avoid contamination the outmost ca. two centimetres of peat was removed. The subsamples were stored in a cold room at 6°C in plastic bags.

### Plant macrofossil analysis

Plant macrofossil analysis was performed at 2-cm resolution. Where prominent changes in plant composition occurred, the resolution was increased to contiguous centimetres. Volumetric samples of 5 cm<sup>3</sup> were carefully rinsed under running water using a 140-µm sieve. From the residue, proportions of the main peat components (such as *Sphagnum*,

non-*Sphagnum* mosses, Cyperaceae remains, wood, roots, and leaves) were estimated as percentages of a total sample volume under a stereomicroscope and a high-power light microscope. Seeds, leaves and charred remains were counted as exact numbers. Plant macrofossil analysis followed Mauquoy and van Geel (2007), modified by Välranta *et al.* (2007). If plant remains were unidentifiable to plant type level, proportion of unidentified organic matter (UOM) was estimated. Identification followed e.g. Laine *et al.* (2009), Mauquoy and van Geel (2007), and Euroala *et al.* (1992). A reference collection was also available. Software C2 (v. 1.7.7) (Juggins, 2007) and Tilia 2.0.41 were used to produce the diagrams.

### Chronology

Radiocarbon ( $^{14}\text{C}$ ) accelerator mass spectrometry (AMS) and lead ( $^{210}\text{Pb}$ ) dating methods were combined to establish accurate chronologies and build age-depth models (van Der Plicht, 2004; Ali *et al.*, 2008). In total 16 samples were sent to Poznan Radiocarbon Laboratory (Poznan, Poland) and three samples to the Finnish Museum of Natural History (LUOMUS, Helsinki, Finland) for  $^{14}\text{C}$  dating (Table S1). Bulk peat samples, cleaned of roots and rootlets when possible (Holmquist *et al.*, 2016), were used for  $^{14}\text{C}$  dating. Samples from the base of each peat monolith, layers representing vegetation changes and/or intermediate layers (LG2.2 and K2.3) were selected for dating.


To create accurate chronologies for recent decades,  $^{210}\text{Pb}$  dating was performed at the University of Exeter, UK, using alpha-spectrometry.  $^{210}\text{Pb}$  was measured at 2-cm intervals for each full-length monolith. Freeze-dried subsamples of 0.14–0.51 g were analysed for  $^{210}\text{Pb}$  activity after spiking them with a  $^{209}\text{Po}$  yield tracer, following the methods described by Kelly *et al.* (2017) and Estop-Aragonés *et al.* (2018).  $^{210}\text{Pb}$  ages were obtained using the Constant Rate of Supply model (CRS) (Appleby and Oldfield, 1978). Age-depth models for each core (Fig. 2) were created with BACON v2.3.3 package in R version 3.4.3 (R Core Team, 2016). BACON divides the dated cores into sections and applies Bayesian statistics with prior information to reconstruct the accumulation, providing weighted mean ages (Blaauw, 2010; Blaauw and Christen, 2011) that are further used for calculations without chronological error ranges.  $^{14}\text{C}$  ages were internally calibrated using the INTCAL 13 calibration curve (Reimer, 2013) and modern dates (pMC % modern carbon) were converted to radiocarbon ages applying NH Zone 1 post bomb curve (Hua *et al.*, 2013).

### Carbon accumulation rates ACAR and RERCA values

For peat and carbon accumulation calculations, dry bulk density ( $\text{g}/\text{cm}^3$ ) was measured contiguously for every cm after freeze-drying volumetric subsamples of  $5\text{ cm}^3$  and by dividing the dry mass (g) by the peat fresh volume ( $\text{cm}^3$ ). Carbon and nitrogen (C/N) content measurements were performed at 4-cm intervals, at the University of Helsinki, using a LECO TruSpec micro Elemental Determinator and these results were applied to calculate average carbon values.


1  
2  
3 148 To estimate the temporal variations in apparent carbon accumulation rates (ACAR, g C m<sup>-2</sup> yr<sup>-1</sup>), the carbon mass of  
4  
5 149 every 1-cm increment (g/m<sup>3</sup>) was multiplied by the corresponding vertical peat accumulation rate (m/yr.) (Turunen *et*  
6  
7 150 *al.*, 2002) based on the age-depth models. Average, non-cumulative, apparent recent rates of carbon accumulation  
8  
9 151 (RERCA) (cf. Turunen 2003) were calculated for three periods; post AD 1950 until present, post AD 1900 until present,  
10  
11 152 and post AD 1850 until present, following a procedure introduced in Lamarre *et al.* (2012). These periods mainly  
12  
13 153 represent incompletely decomposed (acrotelm) peat and are not straight forwardly comparable with results yielded from  
14  
15 154 older highly decomposed, water saturated, and anoxic (catotelm) peat.  
16  
17 155 **Results**  
18  
19  
20 156 *Chronology and peat accumulation*  
21  
22 157 Age-depth models show that peat accumulation has not been constant over the accumulation history and ages at the  
23  
24 158 base of the monoliths vary from ca. 2500 cal. BP (K1P2) to ca. cal. AD 1960 (K2.3 and LG2.2) (Fig. 2). However, in  
25  
26 159 this study, we focus on the period of the last ca. 200 years. Mean peat accumulation rates range from 1.6 mm/yr to 5.4  
27  
28 160 mm/yr (Table 2). On average, accumulation rates increase towards the present, starting from the 1950s (Figs. 3–6).  
29  
30 161 Within both regions, peatland specific variation in accumulation rates exists (Table 2). The K1P2 age-depth model  
31  
32 162 suggests a slowdown in peat accumulation between ca. 1800 cal. BP and 95 cal. BP (19-15 cm) when the peat  
33  
34 163 accumulation rate is only 0.02 mm/yr. Alternatively either of the <sup>14</sup>C ages is an outlier. <sup>210</sup>Pb activity ceases already at  
35  
36 164 13 cm below the surface in K1P2 (Fig. 2). For cores K2.3 and LG2.2 (both 33 cm in length), the <sup>210</sup>Pb analyses did not  
37  
38 165 reach the zero activity level (Table S2). In order to establish age-depth models for these two cores, we <sup>14</sup>C-dated  
39  
40 166 additional peat samples (Table S1 and Fig. 2). Additionally, the <sup>14</sup>C basal-age of the record LG2.3 was younger than the  
41  
42 167 corresponding age suggested by the <sup>210</sup>Pb chronology (Table S2) and the BACON age-depth model therefore excluded  
43  
44 168 the <sup>14</sup>C age as an outlier (Fig. 2). A charcoal layer with charred plant remains in LG2.2 was dated to ca. cal. AD 1940  
45  
46 169 (21 cm) which corresponds to a previously reported fire in AD 1941 (SOPFEU, 2004) and supports the reliability of the  
47 170 chronology (Figs 2, 5).  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60





**Figure 2.** BACON age-depth models.  $^{210}\text{Pb}$  age-ranges are in green (for ★LG2.2 and ★K2.3 green indicates the coring year), and  $^{14}\text{C}$  dates (cal. BP) are in blue. The grey shading indicates the possible age-range and darkest grey demarks most likely range. The thin red line in the middle is the weighted mean age based on the model.

1  
2  
3 176 The average core-specific bulk density (BD) with standard deviation ( $\pm$ SD) varies between  $0.05\pm0.01$  g/cm<sup>3</sup> (LG2.3)  
4  
5 177 and  $0.11\pm0.04$  g/cm<sup>3</sup> (K1P2) with an increasing trend towards older peat layers (Figs. 3–6). The mean BD of all the  
6  
7 178 cores was  $0.07\pm0.03$  g/cm<sup>3</sup>. Carbon content analysis yielded an average of  $45.5\pm1.7\%$ . Carbon averages calculated over  
8  
9 179 an individual peat core ranged from  $43.9\pm1.4\%$  (K2.3) to  $47.1\pm2.7\%$  (K1P2). C/N ratio varied between 14.3 and 62.8  
10  
11 180 with an average of  $36.4\pm11.9$ . The overall trend in C/N values increased towards the surface (Figs. 3–6).  
12  
13 181 *Past vegetation succession*  
14  
15  
16 182 *Kuujuarapik (K1P)*  
17  
18 183 At first, until ca. cal. AD 1710 (33 cm), *S. fuscum* dominates in K1P1, after which the dry assemblage is replaced by  
19  
20 184 wet *S. lindbergii* and Cyperaceae (Fig. S1). Between ca. cal. AD 1960 and cal. AD 2000, at 13–7 cm, the K1P1 record  
21  
22 185 shows a change from a wet assemblage back to a drier assemblage with *Dicranum* sp. and dwarf shrubs. From ca. cal.  
23  
24 186 AD 1990 (9 cm), a *S. fuscum*-ericaceous assemblage dominates.  
25  
26 187 K1P2 is first dominated by *S. fuscum* with numerous *Picea* sp. needles (33–19 cm), woody material and sparse charred  
27  
28 188 fragments (33–12 cm). Cyperaceous taxa are most abundant between 20–11 cm up to ca. cal. AD 1950. Wet *Sphagnum*  
29  
30 189 sect. Cuspidata was identified at 13–11 cm. Shrub roots replace the *Sphagnum* dominance between ca. cal. AD 1950 and  
31  
32 190 1995 (11–7 cm). *S. fuscum* dominates the vegetation from ca. cal. AD 1995 (7–0 cm) to present.  
33  
34 191 In K1P3, Cyperaceae and *Sphagnum* assemblages alternate. First (from 31 cm, ca. cal. AD 1765), Cyperaceae spp.  
35  
36 192 dominate until ca. cal. AD 1780 (30–29 cm) when the assemblage shifts to *S. lindbergii*. Non-*Sphagnum* bryophytes e.g.  
37  
38 193 *Straminergon stramineum* and *Exannulata* group species are present through these phases. After ca. cal. AD 1930 (21  
39  
40 194 cm) *S. lindbergii* is first accompanied by, and then in ca. cal. AD 1970 replaced with Cyperaceae species.  
41  
42 195 *Trichophorum cespitosum* prevails with dwarf shrubs between ca. cal. AD 1970 and ca. cal. AD 1990 (17–15 cm). After  
43  
44 196 ca. cal. AD 1990 (14 cm), *S. russowii* re-establishes the *Sphagnum* assemblage with small quantities of *Trichophorum*  
45  
46 197 *cespitosum* and ericaceous plants.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




**Figure 3.** Plant composition presented as main vegetation types with different colours (%). Y-axis indicates the AD year from 1800 until the coring year 2017. Symbol // is used to indicate that the record continues to older ages. Continuous line represents K1P1 record, dashed line K1P2, and dotted line K1P3. On the top x-axis  $\text{g C m}^{-2} \text{yr}^{-1}$  stands for ACAR, mm/yr for peat accumulation and BD ( $\text{g/cm}^3$ ) for bulk density. RW zone (recent warming) and LIA zone (Little Ice Age) are shown with grey horizontal lines.

#### Kuujuarapik (K2)

Cyperaceae spp. dominate the K2.1 record from ca. cal. AD 1825 until ca. cal. AD 1845 (29 cm), when the plant assemblage changes to *Sphagnum fuscum* dominance (Fig. 4). At ca. cal. AD 1995 (13 cm), slightly more Cyperaceae remains, dwarf shrub roots, and Ericaceae leaves (mainly *Chamaedaphne calyculata*) occur.

In K2.2, macroscopic charcoal is detected between ca. cal. AD 1900 (31 cm) and ca. cal. AD 1935 (27 cm). Cyperaceae spp. are present until ca. cal. AD 1980 (21 cm) but are replaced by *S. fuscum* via *S. sect. subsecunda* and ericaceous roots. *S. fuscum* dominates the plant assemblages until the present-day together with *Trichophorum cespitosum* and Ericaceae (Fig. 4, S1).

In K2.3, the assemblages of cyperaceous plants and *S. fuscum* vary between ca. cal. AD 1960 and 1990 (33-16 cm) (Fig. 4, S1). *Chamaedaphne calyculata* seeds are numerous around cal. AD 1980s (21-18 cm) and *Myrica anomala* is abundant (up to 40 %) between ca. cal. AD 1985 and ca. cal. AD 1990 (18-16 cm). From ca. cal. AD 1990 to the present-day, *S. fuscum* dominates the plant assemblages. Dwarf shrub roots, *Trichophorum cespitosum*, and *Polytrichum strictum* occur throughout the record in low quantities.



**Figure 4.** Plant composition presented as main vegetation types with different colours (%). Y-axis indicates the AD year from 1800 until the coring year 2017. Symbol \* indicates where charcoal and/or charred plants were found. Continuous line represents K2.1 record, dashed line K2.2, and dotted line K2.3. On the top x-axis  $\text{g C m}^{-2} \text{ yr}^{-1}$  stands for ACAR, mm/yr for peat accumulation and BD ( $\text{g/cm}^3$ ) for bulk density. RW zone (recent warming) and LIA zone (Little Ice Age) are shown with grey horizontal lines.


#### Radisson (LG2)

The LG2.1 vegetation assemblage is predominantly composed of *Sphagnum capillifolium* together with ligneous plants and some lichens from ca. cal. AD 1850 until ca. cal. AD 1945 (20 cm) (Fig. 5, S1). A layer with charred plant remains occurred ca. cal. AD 1940 (21 cm). After the fire to the present-day, *Sphagnum fuscum* dominates the plant assemblage occasionally accompanied by *Myrica anomala*.

No substantial changes are detected in the LG2.2 plant assemblages. At first, *S. fuscum* is occurring together with cf. *S. subfulvum*. After ca. cal. AD 1965, cyperaceous remains become more abundant. *S. fuscum* becomes dominant again after ca. cal. AD 1970 (27 cm), together with some sparse cyperaceous taxa and dwarf shrub roots that prevail throughout the record (Fig. 5, S1).

In LG2.3, charcoal and charred plant fragments are identified at ca. cal. AD 1850 (31 cm). Until ca. cal. AD 1885 (28 cm), *S. balticum* prevails with *S. fuscum*, *S. rubellum*, and *Trichophorum cespitosum*. After ca. cal. AD 1885, *S. balticum* disappears and *S. fuscum* dominates the plant assemblages with *S. rubellum*, until present. Dwarf shrub roots

and *Myrica anomala* are scarce throughout. Ericaceous leaves become more abundant after ca. cal. AD 1965 (19 cm) (Fig. 5, S1).



**Figure 5.** Plant composition presented as main vegetation types with different colours (%). Y-axis indicates the AD year from 1800 until the coring year 2017. Symbol \* indicates where charcoal and/or charred plants were found. Continuous line represents LG2.1 record, dashed line LG2.2, and dotted line LG2.3. On the top x-axis g C m<sup>-2</sup> yr<sup>-1</sup> stands for ACAR, mm/yr for peat accumulation and BD (g/cm<sup>3</sup>) for bulk density. RW zone (recent warming) and LIA zone (Little Ice Age) are shown with grey horizontal lines.


#### Radisson (Rad)

In Rad1 (from 38 cm, ca. cal. AD 1840), *S. fuscum* dominates the assemblage accompanied by *S. rubellum* and *S. capillifolium*. Dwarf shrub roots are abundant (up to 40%) and lichens are found from ca. cal. AD 1920 until ca. cal. AD 1960 (29-23 cm). Charcoal is found between cal. AD 1920s and 1950s (29-25 cm) (Fig. 6, S1). After the charcoal layers, *S. balticum* and *Myrica anomala* are found and ericaceous plants become more abundant. Towards present, the prevalence of roots declines and *Sphagnum* represents up to 80% of plant remains.

In Rad2 (37 cm: from ca. cal. AD 1770), *S. fuscum* dominates the plant assemblage throughout the record, with a reduction from 75-90% to 50% after cal. AD 1970s (18-16 cm). Charcoal and charred plant fragments occur between late cal. AD 1960s and mid-1970s (21-17 cm). Some lichens are found underneath the charcoal layer until the cal. AD 1930s (23 cm) and at 16 cm which corresponds to the late cal. AD 1980s. Cyperaceous taxa are found in small



1  
2  
3 253 abundance until ca. cal. AD 1990 (13 cm). In the plant assemblages following the charcoal layers, *Myrica anomala*  
4  
5 254 appears with ericaceous shrubs up to the present-day surface.  
6  
7 255 From the base of Rad3 record (36 cm, ca. cal. AD 1780), until ca. cal. AD 1820 (32 cm), cf. *S. subfulvum* is abundant  
8  
9 256 together with Cyperaceae species. Charcoal and charred plant fragments are detected between ca. cal. AD 1820-1890  
10  
11 257 (32-24 cm). After ca. cal. AD 1820, *Sphagnum* remains diminish with an increase in dwarf shrub roots. *Sphagnum* sect.  
12  
13 258 Cuspidata occurs until ca. cal. AD 1880 (25 cm), when replaced by *S. fuscum* and *S. rubellum*. *Picea mariana* needles  
14  
15 259 are numerous around ca. cal. AD 1880s (25 cm). After the last charcoal layer (ca. cal. AD 1890), *S. rubellum* and *S.*  
16  
17 260 *fuscum* accompanied by dwarf shrub roots dominate the record (Fig. 6, S1).



41 262 **Figure 6.** Plant composition presented as main vegetation types with different colours (%). Y-axis indicates the AD  
42  
43 263 year from 1800 until the coring year 2017. Symbol // is used to indicate that the record continues to older ages, and  
44  
45 264 symbol \* indicates where charcoal and/or charred plants were found. Continuous line represents Rad1 record, dashed  
46  
47 265 line Rad2, and dotted line Rad3. On the top x-axis g C m<sup>-2</sup> yr<sup>-1</sup> stands for ACAR, mm/yr for peat accumulation and BD  
48  
49 266 (g/cm<sup>3</sup>) for bulk density. RW zone (recent warming) and LIA zone (Little Ice Age) are shown with grey horizontal lines.

51  
52 267 *ACAR and RERCA variability*  
53  
54 268 Because the ages at the base of the monoliths vary, the oldest dated to 2510 cal. BP (K1P2) and youngest to -10 cal. BP  
55  
56 269 (cal. AD 1960, K2.3 and LG2.2), we focused on three defined RERCA periods: from AD 1850, from AD 1900, and  
57  
58 270 from AD 1950 to the present-day (Table 2). This approach enables us to identify and compare possible changes in  
59  
60 271 recent carbon accumulation rates. In general, the post AD 1950 RERCA rates are the highest (Table 2, Figs. 3–6), partly


due to incomplete decomposition. At K2, RERCA values were consistently high (Table 2). The lowest post AD 1950 RERCA rates were detected for the Radisson region peatlands: LG2.1: 65.3 g C m<sup>-2</sup> yr<sup>-1</sup> and Rad3: 50.7 g C m<sup>-2</sup> yr<sup>-1</sup>. However, for these same peatlands, much higher rates were also derived: LG2.2: 136.9 g C m<sup>-2</sup> yr<sup>-1</sup> and Rad1: 103.3 g C m<sup>-2</sup> yr<sup>-1</sup>.

**Table 2.** Core length (cm) represents the thickness of the unfrozen peat layer at the time of the sampling. Water table depth (WTD). Average peat accumulation rates (peat accu.) mm/yr. Average recent apparent carbon accumulation rates (RERCA) (g C m<sup>-2</sup> yr<sup>-1</sup>) were calculated for periods AD 1850, AD 1900, and AD 1950 until present. NA: not available indicates that the chronology does not reach that far back in time. Basal ages from the core bottoms (cal. BP) are rounded to nearest five.

Core	Core length (cm)	WTD (cm)	Peat accu. (mm/yr.)	Basal age of the core Cal. BP	RERCA AD1950	RERCA AD1900	RERCA AD1850
K1P1	39	13	1.7	340	70.1	60.1	53.0
K1P2	35	18	1.6	2530	91.7	82.6	74.3
K1P3	31	18	3.2	185	121.0	108.4	101.8
K2.1	32	27	3.5	140	102.1	89.3	83.3
K2.2	31	22	4.9	50	149.1	140.6	NA
K2.3	33	20	5.4	-10	147.9	NA	NA
LG2.1	34	20	2.3	115	65.3	64.6	63.1
LG2.2	33	21	5.4	-10	136.9	NA	NA
LG2.3	34	18	3.2	120	91.9	77.7	71.1
Rad1	38	23	3.2	120	103.3	95.7	89.6
Rad2	37	14	2.9	190	100.0	89.4	81.9
Rad3	36	19	1.7	180	50.7	51.0	54.1

ACAR variability is high (Fig. 7) and mainly varies with depth, with higher rates towards the peat surfaces (Figs. 3–6). RERCA ranges and exact values for the three focus periods are higher for Kuujjuarapik than for Radisson peatlands (e.g. AD 1950 RERCAs are 24% higher in Kuujjuarapik region than in Radisson) (Fig. 7). The site-combined averages for the three periods were consistently higher in Kuujjuarapik than in Radisson, but also the lowest apparent carbon accumulation rates were calculated for Kuujjuarapik (Fig. 7).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17 287  
18 288  
19  
20 289  
21  
22 290  
23  
24  
25 291  
26  
27 292  
28  
29 293  
30  
31 294  
32  
33 295  
34  
35 296  
36  
37 297  
38  
39 298  
40  
41 299  
42  
43 300  
44  
45 301  
46  
47 302  
48  
49 303  
50  
51 304  
52 305  
53  
54 306  
55  
56  
57 307  
58  
59  
60



**Figure 7.** ACAR (g C m<sup>-2</sup> yr<sup>-1</sup>) variability for each core (left panel) and combined RERCA (g C m<sup>-2</sup> yr<sup>-1</sup>) variability for the two study regions, Kuujjuarapik and Radisson, focusing on three periods: from AD 1950, from AD 1900, and from AD 1850 to present (right panel). See legend in the left panel.

**Discussion**

Variations in vegetation succession patterns between and within sites neither fully support nor reject our first research question that plant assemblages are changing synchronously in response to rapid recent warming. In contrast to our second research question, the northern Kuujjuarapik peatlands had higher RERCA rates when compared to the more southern Radisson site but supporting the research question, carbon accumulation rates increased towards recent decades. This increase is, however, partly expected due to the incomplete state of decomposition even if the climate had not become warmer. Plant assemblages of K2, Rad, and LG2 peatlands revealed asynchronous response to post-LIA warming (Fig. 4–6). Moreover, all sampling sites indicated more recent decadal-scale changes; yet the exact timing varies (Figs. 3–6). In K1P peatland, the vegetation change was synchronous for all records starting from ca. cal. AD 1950s and lasting until mid-1990s resulting with an establishment of a drier *Sphagnum* lawn microform. In turn, at K2 peatland, the vegetation assemblages showed successional changes around cal. AD 1970s (K2.2), 1980s (K2.3), and 1990s (K2.1). However, K2 peatland has different permafrost dynamics and microtopography gradient from K1P palsa peatland. Vegetation community changes in Rad peatland and the LG2.1 record around cal. AD 1940s might have been fire-induced as suggested by the charred plant remains and charcoal in the peat records. Kuujjuarapik vegetation assemblages seem to have experienced more synchronous and distinct changes than the southern, Radisson assemblages. This could be linked to changes in thawing permafrost dynamics in the northern Kuujjuarapik site.

*Peat properties and peat and carbon accumulation*



Our BD values corresponded to previous studies from Northern Ontario where BD ranged from 0.0034 to 0.62 g/cm<sup>3</sup> with an average of 0.093±0.041 g/cm<sup>3</sup> (Holmquist et al., 2014). Our average carbon content was slightly lower than the 50% which is often used for peat carbon content calculations for boreal and subarctic to Arctic regions (Turunen et al., 2002; Treat et al., 2016), but close to an average reported for northern peatlands 46.8±6.1% (Loisel et al., 2014). Our C/N averages are comparable with other studies, but our lowest C/N values were lower than those reported by Treat et al. (2016) (C/N 30-62).

Previous results from Kuujjuarapik have shown a notable rise of peat accumulation rates since the beginning of the 20th century, with average RERCAs: 133-147 g C m<sup>-2</sup> yr<sup>-1</sup> from cal. AD 1950 to present, 73-81 g C m<sup>-2</sup> yr<sup>-1</sup> from cal. AD 1900 to present, and 52-62 g C m<sup>-2</sup> yr<sup>-1</sup> from cal. AD 1850 to present (Lamarre et al., 2012). The highest individual RERCA values of Lamarre et al. (2012) exceeded 160 g C m<sup>-2</sup> yr<sup>-1</sup> and were interpreted to reflect the recent increase in annual temperatures.

High peat accumulation rates of 0.5 cm/yr, with an average of 0.037 cm/yr, and high RERCA values (from cal. AD 1850) between 52.8 and 114.5 g C m<sup>-2</sup> yr<sup>-1</sup> with a mean of 73.6 g C m<sup>-2</sup> yr<sup>-1</sup>, have been reported for Eastmain, James Bay area south from Radisson (Loisel and Garneau, 2010). Moreover, a mean carbon accumulation rate over the past ca. 300 years of 56.4 g C m<sup>-2</sup> yr<sup>-1</sup>, has been obtained from the same region (van Bellen et al., 2011b). Turunen et al. (2004) reported a carbon accumulation rate of 73 g C m<sup>-2</sup> yr<sup>-1</sup> for eastern Canadian bogs, covering the last 150 years. High carbon accumulation rates of > 150 g C m<sup>-2</sup> yr<sup>-1</sup> over the past ca. 200 years in northeastern maritime Québec have been put forward by Magnan and Garneau (2014a). In contrast, Hudson Bay Lowland records from Northern Ontario indicated no rise in carbon accumulation rates for the recent decades, i.e. less than 30 g C m<sup>-2</sup> yr<sup>-1</sup> (Bunbury et al., 2012). There, the lead activity covered only the topmost 6 cm, while in the current study, where the core lengths were between 31 cm and 39 cm, the zero <sup>210</sup>Pb activity level was sometimes not reached at all (K2.3 and LG2.2), which creates uncertainty for the chronologies. However, in K1P2, the lead activity covered only the topmost 13 cm, resembling the results of Bunbury et al. (2012). These previous studies were conducted in peatlands resembling ours and samples were mainly collected from similar microhabitats. It has been suggested that peat and carbon accumulation rates are fast for *Sphagnum*-dominated habitats under high effective moisture conditions and slower when sedges dominate in warm and dry conditions (Nichols et al. 2014). In the La Grande region, a rise in peat accumulation rates together with a recent change towards *Sphagnum*-dominated plant communities has been detected (Beaulieu-Audy et al., 2009, Pratte et al. 2017). In our study, however, we found no correlation between *Sphagnum* assemblages and carbon accumulation (Pearson correlation: r=0.09, p=0.21).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

337 *Drivers behind accumulation patterns*

338 Intermediate peatland microforms, selected for this study, have higher vertical peat and carbon accumulation rates than,  
339 for instance, wet hollows or high and dry hummocks (Belyea and Clymo, 2001). *Sphagnum* establishment promotes  
340 rapid peat accumulation and hence peatland vegetation community structure strongly influences carbon sequestration  
341 (Tuittila *et al.*, 2012; Loisel and Yu, 2013). However, it is often a challenge to separate internal and site-specific and  
342 external, e.g. climate driven, forcing (Tuittila *et al.*, 2007, 2012; Loisel and Garneau, 2010; van Bellen *et al.*, 2011b;  
343 Loisel and Yu, 2013). Changes in hydrology and vegetation can also be caused by long-term hydrosereal succession  
344 from fen to bog (ombrotrophication) that is driven by both internal and external forcing (Yu *et al.* 2009; Väiranta *et al.*,  
345 2017).

346 PAR0 and growing season length have been connected to *Sphagnum* productivity (NPP) (Loisel *et al.*, 2012; Gallego-  
347 Sala *et al.*, 2018). Accordingly, peat accumulation should be positively correlated to GDD0 (Clymo *et al.*, 1998;  
348 Charman *et al.*, 2013; Holmquist *et al.*, 2014). In addition, moisture is an important factor controlling carbon  
349 accumulation and peatland dynamics (Swindles *et al.*, 2015; Galka *et al.*, 2017; Zhang *et al.*, 2018a). Both precipitation  
350 and GDD0 have increased since the 1960s and 1970s in our study sites (Environment Canada, 2018). In the Hudson  
351 Bay region, sea ice cover has decreased up to  $11.3\% \pm 2.6\%$  per decade from AD 1971 to AD 2008, which is affecting  
352 moisture and heat transfer patterns of the area (Tivy *et al.*, 2011) and may also affect peat accumulation processes.

353 Our aim was to study recent peat and carbon accumulation patterns by collecting peat records from uniform lawn  
354 *Sphagnum* microforms. Because establishment of robust chronologies for young peat sections is challenging (see  
355 however e.g. Goslar *et al.* 2005), in this study we combined two dating methods that support each other (e.g. Turetsky *et*  
356 al. 2004). Our results indicated high accumulation rates during recent decades. Previous studies focussing on both the  
357 last millennium (Charman *et al.*, 2013; Garneau *et al.*, 2014; Loisel *et al.*, 2014) and recent decades (Zhang *et al.*,  
358 2018b) showed that relatively high accumulation rates are likely caused by increased carbon inputs rather than by  
359 reduced decomposition. We acknowledge the difficulty of interpreting the recent peat accumulation rates due to the  
360 incomplete decomposition process and lower compaction of the surface peat and restraints on the chronologies. Despite  
361 these inaccuracies, by using specified focus periods, the data nevertheless allow spatio-temporal comparisons and the  
362 assessment of data against the previously published data carried out using similar protocols.

363 *Future implications*

364 Observed and anticipated warmer and wetter climate conditions for the coming decades (Kirtman *et al.*, 2013,  
365 Environmental Canada, 2018) should be beneficial for peat accumulation (Payette *et al.*, 2004). Following this, a

potential increase of carbon storage has been suggested (Holmquist *et al.*, 2014; Gallego-Sala *et al.* 2018). However, contrasting views also exist where modelling exercises suggest that in eastern Canada, peatlands will turn into a carbon source due to increased decomposition under warmer climate, despite the potential increase in carbon sequestration (Chaudhary *et al.*, 2017). McGuire *et al.* (2018) modelled the vegetation response to climate warming and CO<sub>2</sub> fertilization under different warming scenarios. They predicted that depending on the magnitude of warming, northern permafrost soils could act as a net sink of carbon, but if the warming is more prominent, substantial soil carbon losses could appear after 2100 (McGuire *et al.*, 2018). In some of the projections, vegetation was largely responsible for the carbon intake, but the other projections indicated that vegetation carbon intake could not compensate the losses (McGuire *et al.*, 2018). To address the question how high-latitude peatland vegetation and carbon dynamics will respond and have responded to recent warming, more data are needed including high-resolution chronologies and multiple cores, as presented in this study. The presented data highlight the importance of studying multiple peat sections from one study site and preferably the approach should be extended to regional-scales as stated by our third research question and the internal variability needs to be taken into account when thinking about upscaling and modelling of results (Loisel and Garneau, 2010; Lamarre *et al.*, 2012; Mathijssen *et al.*, 2017; Zhang *et al.*, 2018a).

## Conclusions

This study analysed twelve shallow surface peat cores for plant macrofossils and carbon accumulation to reveal recent changes in habitat conditions and peat properties in northern peatlands of eastern Canada. Our results indicated more prominent changes in vegetation dynamics and carbon accumulation in the northernmost study sites than in the southern sites. These changes may be linked to larger increase in the growing season length, different climatic conditions, and changing permafrost dynamics of the northern region. High-resolution chronologies, by applying multiple complementary dating methods, is essential to reveal the most recent changes. Our study shows that in order to exclude the influence of internal peatland variability on detected changes in accumulation patterns, multiple cores are needed to capture the genuine regional-scale climate signal.

## Acknowledgements

We thank N. Chaumont for fieldwork assistance, N. Sanderson for helping with establishing the chronologies, and R. Gauthier for rechecking one *Sphagnum* species. The Academy of Finland (code 296519) and Natural Sciences and Engineering Research Council of Canada (NSERC; No. 250287) discovery grant to M.G funded this project. We thank the three anonymous reviewers for their constructive comments.

## References

- 1  
2
- 3 395 Abram, N.J. et al. and the PAGES 2k Consortium (2016) 'Early onset of industrial-era warming across the oceans and  
4 396 continents', *Nature*, 536, pp. 411–418, doi:10.1038/nature19082.
- 5  
6 397 Ali, A. A. *et al.* (2008) 'Recent peat accumulation rates in minerotrophic peatlands of the Bay James region, Eastern  
7 398 Canada, inferred by  $^{210}\text{Pb}$  and  $^{137}\text{Cs}$  radiometric techniques', *Applied Radiation and Isotopes*, 66(10), pp. 1350–1358.  
8 399 doi: 10.1016/j.apradiso.2008.02.091.
- 9 400 Allard, M. and K.-Seguin, M. (1987) 'Le pergélisol au Québec nordique : bilan et perspectives', *Géographie physique  
10 401 et Quaternaire*, 41(1), p. 141. doi: 10.7202/032671ar.
- 11  
12 402 Appleby, P. G. and Oldfield, F. (1978) 'The calculation of lead-210 dates assuming a constant rate of supply of  
13 403 unsupported  $^{210}\text{Pb}$  to the sediment', *Catena*, 5, pp. 1–8.
- 14 404 Arlen-Pouliot, Y. and Bhiry, N. (2005) 'Palaeoecology of a palsa and a filled thermokarst pond in a permafrost peatland,  
15 405 subarctic Québec, Canada', *Holocene*, 15(3), pp. 408–419. doi: 10.1191/0959683605hl818rp.
- 16  
17 406 Arseneault, D. and Sirois, L. (2004) 'The millennial dynamics of a boreal forest stand from', *Journal of Ecology*, 92, pp.  
18 407 490–504.
- 19 408 Beaulieu-Audy, V. *et al.* (2009) 'Holocene palaeoecological reconstruction of three boreal peatlands in the la Grande  
20 409 Rivière region, Québec, Canada', *Holocene*, 19(3), pp. 459–476. doi: 10.1177/0959683608101395.
- 21  
22 410 van Bellen, S., Garneau, M. and Booth, R. K. (2011a) 'Holocene carbon accumulation rates from three ombrotrophic  
23 411 peatlands in boreal Quebec, Canada: Impact of climate-driven ecohydrological change', *Holocene*, 21(8), pp. 1217–  
24 412 1231. doi: 10.1177/0959683611405243.
- 25  
26 413 van Bellen, S. *et al.* (2011b) 'Quantifying spatial and temporal Holocene carbon accumulation in ombrotrophic  
27 414 peatlands of the Eastmain region, Quebec, Canada', *Global Biogeochemical Cycles*, 25(2), pp. 1–15. doi:  
28 415 10.1029/2010GB003877.
- 29 416 Belyea, L. R. and Clymo, R. S. (2001) 'Feedback control of the rate of peat formation', *Proceedings of the Royal  
30 417 Society B: Biological Sciences*, 268(1473), pp. 1315–1321. doi: 10.1098/rspb.2001.1665.
- 31  
32 418 Blaauw, M. (2010) 'Methods and code for "classical" age-modelling of radiocarbon sequences', *Quaternary  
33 419 Geochronology*. Elsevier B.V, 5(5), pp. 512–518. doi: 10.1016/j.quageo.2010.01.002.
- 34 420 Blaauw, M. and Christen, J. A. (2011) 'Flexible paleoclimate age-depth models using an autoregressive gamma  
35 421 process', *Bayesian Analysis*, 6(3), pp. 457–474. doi: 10.1214/11-BA618.
- 36  
37 422 Bunbury, J., Finkelstein, S. A. and Bollmann, J. (2012) 'Holocene hydro-climatic change and effects on carbon  
38 423 accumulation inferred from a peat bog in the Attawapiskat River watershed, Hudson Bay Lowlands, Canada',  
39 424 *Quaternary Research (United States)*. Université du Québec à Montréal, 78(2), pp. 275–284. doi:  
40 425 10.1016/j.yqres.2012.05.013.
- 41 426 Carroll, P. and Crill, P. (1997) 'Carbon balance of a temperate poor fen', *Global Biogeochemical Cycles*, 11(3), pp.  
42 427 349–356. doi: 10.1029/97GB01365.
- 43  
44 428 Charman, D. J. *et al.* (2013) 'Climate-related changes in peatland carbon accumulation during the last millennium',  
45 429 *Biogeosciences*, 10(2), pp. 929–944. doi: 10.5194/bg-10-929-2013.
- 46  
47 430 Charman, D. J., Aravena, R. and Warnert, B. G. (1994) 'Carbon Dynamics in a Forested Peatland in North-Eastern  
48 431 Ontario, Canada' Warner Published by: British Ecological Society, 82(1), pp. 55–62.
- 49 432 Charman, D. J. *et al.* (2015) 'Drivers of Holocene peatland carbon accumulation across a climate gradient in  
50 433 northeastern North America', *Quaternary Science Reviews*. Elsevier Ltd, 121, pp. 110–119. doi:  
51 434 10.1016/j.quascirev.2015.05.012.
- 52  
53 435 Chaudhary, N., Miller, P. A. and Smith, B. (2017) 'Modelling past, present and future peatland carbon accumulation  
54 436 across the pan-Arctic region', *Biogeosciences*, 14(18), pp. 4023–4044. doi: 10.5194/bg-14-4023-2017.
- 55 437 Clymo, A. R. S., Turunen, J. and Tolonen, K. (1998) 'Nordic Society Oikos Carbon Accumulation in Peatland  
56 438 Published by : Wiley on behalf of Nordic Society Oikos Stable URL : <http://www.jstor.org/stable/3547057>  
57 439 REFERENCES Linked references are available on JSTOR for this article : You may need to log in to J', 81(2), pp. 368–  
58 440 388.
- 59  
60 441 Collins, M. *et al.* (2013) 'Long-term Climate Change: Projections, Commitments and Irreversibility', *Climate Change*

- 2013: *The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 1029–1136. doi: 10.1017/CBO9781107415324.024.
- Crowther, T. W. *et al.* (2016) ‘Quantifying global soil carbon losses in response to warming’, *Nature*, 540(7631), pp. 104–108. doi: 10.1038/nature20150.
- Davidson, E. A. and Janssens, I. A. (2006) ‘Temperature sensitivity of soil carbon decomposition and feedbacks to climate change’, *Nature*, 440(7081), pp. 165–173. doi: 10.1038/nature04514.
- Dorrepaal, E. *et al.* (2009) ‘Carbon respiration from subsurface peat accelerated by climate warming in the subarctic’, *Nature*. Nature Publishing Group, 460(7255), pp. 616–619. doi: 10.1038/nature08216.
- Dyke, A. S. and Prest, V. K. (1987) ‘Late Wisconsinan and Holocene History of the Laurentide Ice Sheet’, *Géographie physique et Quaternaire*, 41(2), p. 237. doi: 10.7202/032681ar.
- Environment Canada. (2018). 1981–2010, 1961–1990 and 1971–2000 Climate Normals and Averages. Retrieved on February 28, 2018, from: [http://climate.weather.gc.ca/climate\\_normals/](http://climate.weather.gc.ca/climate_normals/).
- Estop-Aragonés, C. *et al.* (2018) ‘Limited release of previously-frozen C and increased new peat formation after thaw in permafrost peatlands’, *Soil Biology and Biochemistry*. Elsevier, 118(December 2017), pp. 115–129. doi: 10.1016/j.soilbio.2017.12.010.
- Eurola S. Bendiksen K. and Rönkä A. (1992) Suokasviopas. 2. korjattu painos. *Oulanka Reports*. Viramo J. (editor) Oulanka Biological Station, University of Oulu, (11) pp. 205
- Frolking, S. and Roulet, N. T. (2007) ‘Holocene radiative forcing impact of northern peatland carbon accumulation and methane emissions’, *Global Change Biology*, 13(5), pp. 1079–1088. doi: 10.1111/j.1365-2486.2007.01339.x.
- Frolking, S., Talbot, J. and Subin, Z. M. (2014) ‘Exploring the relationship between peatland net carbon balance and apparent carbon accumulation rate at century to millennial time scales’, *Holocene*, 24(9), pp. 1167–1173. doi: 10.1177/0959683614538078.
- Gałka, M. *et al.* (2017) ‘Vegetation Succession, Carbon Accumulation and Hydrological Change in Subarctic Peatlands, Abisko, Northern Sweden’, *Permafrost and Periglacial Processes*, 28(4), pp. 589–604. doi: 10.1002/ppp.1945.
- Gałka, M. *et al.* (2018) ‘Response of plant communities to climate change during the late Holocene: Palaeoecological insights from peatlands in the Alaskan Arctic’, *Ecological Indicators*, 85(November), pp. 525–536. doi: 10.1016/j.ecolind.2017.10.062.
- Gallego-Sala, A. V *et al.* (2018) ‘Latitudinal limits to the predicted increase of the peatland carbon sink with warming’, *Nature Climate Change*. doi: 10.1038/s41558-018-0271-1.
- Garneau, M. *et al.* (2014) ‘Holocene carbon dynamics of boreal and subarctic peatlands from Québec, Canada’, *Holocene*, 24(9), pp. 1043–1053. doi: 10.1177/0959683614538076.
- Gorham, E. (1991) ‘Northern Peatlands : Role in the Carbon Cycle and Probable Responses to Climatic Warming’, *Ecological Applications*, 1(2), pp. 182–195. doi: 10.2307/1941811.
- Goslar, T. *et al.* (2005) ‘Radiocarbon dating of modern peat profiles: pre- and post-bomb  $^{14}\text{C}$  variations in the construction of age-depth models’, *RADIOCARBON*, Vol 47, Nr 1, p 115–134.
- Holmquist, J. R. *et al.* (2016) ‘A comparison of radiocarbon ages derived from bulk peat and selected plant macrofossils in basal peat cores from circum-arctic peatlands’, *Quaternary Geochronology*, 31. doi: 10.1016/j.quageo.2015.10.003.
- Holmquist, J. R., MacDonald, G. M. and Gallego-Sala, A. (2014) ‘Peatland Initiation, Carbon Accumulation, and 2 ka Depth in the James Bay Lowland and Adjacent Regions’, *Arctic, Antarctic, and Alpine Research*, 46(1), pp. 19–39. doi: 10.1657/1938-4246-46.1.19.
- Hua, Q., Barbetti, M. and Rakowski, A. Z. (2013) ‘Atmospheric Radiocarbon for the Period 1950–2010’, *Radiocarbon*, 55(04), pp. 2059–2072. doi: 10.2458/azu\_js\_rc.v55i2.16177.
- IPCC (2013) ‘Summary for Policymakers’, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, p. 33. doi: 10.1017/CBO9781107415324.
- Ise, T. *et al.* (2008) ‘High sensitivity of peat decomposition to climate change through water-table feedback’, *Nature*

- 1  
2  
3 488 *Geoscience*, 1(11), pp. 763–766. doi: 10.1038/ngeo331.
- 4  
5 489 Jean, M. and Payette, S. (2014) ‘Effect of vegetation cover on the ground thermal regime of wooded and non-wooded  
6 490 palsas’, *Permafrost and Periglacial Processes*, 25(4), pp. 281–294. doi: 10.1002/ppp.1817.
- 7 491 Juggins, S. (2007) ‘C2 user guide: Software for ecological and palaeoecological data analysis and visualization’,  
8 492 *University of Newcastle, Newcastle upon Tyne, UK*, pp. 1–73. Available at:  
9 493 [http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:C2+Software+for+ecological+and+palaeoecological](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:C2+Software+for+ecological+and+palaeoecological+data+analysis+and+visualisation#0)  
10 494 [+data+analysis+and+visualisation#0](http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:C2+Software+for+ecological+and+palaeoecological+data+analysis+and+visualisation#0).
- 11  
12 495 Kelly, T. J. *et al.* (2017) ‘The vegetation history of an Amazonian domed peatland’, *Palaeogeography,*  
13 496 *Palaeoclimatology, Palaeoecology*. Elsevier B.V., 468, pp. 129–141. doi: 10.1016/j.palaeo.2016.11.039.
- 14 497 Kirtman, B. *et al.* (2013) ‘Near-term Climate Change: Projections and Predictability’, *Climate Change 2013 - The*  
15 498 *Physical Science Basis*, pp. 953–1028. doi: 10.1017/CBO9781107415324.023.
- 16  
17 499 Klein, E. S., Yu, Z. and Booth, R. K. (2013) ‘Recent increase in peatland carbon accumulation in a thermokarst lake  
18 500 basin in Southwestern Alaska’, *Palaeogeography, Palaeoclimatology, Palaeoecology*. Elsevier B.V., 392, pp. 186–195.  
19 501 doi: 10.1016/j.palaeo.2013.09.009.
- 20 502 Laine J., Harju P., Timonen T., Laine A., Tuittila E-S., Minkinen K. and Vasander H. (2009) The Intricate Beauty of  
21 503 Sphagnum Mosses – a Finnish Guide to Identification. University of Helsinki Department of Forest Ecology,  
22 504 Publications 39.
- 23  
24 505 Lamarre, A., Garneau, M. and Asnong, H. (2012) ‘Holocene paleohydrological reconstruction and carbon accumulation  
25 506 of a permafrost peatland using testate amoeba and macrofossil analyses, Kuujuaupik, subarctic Qu??bec, Canada’,  
26 507 *Review of Palaeobotany and Palynology*. Elsevier B.V., 186, pp. 131–141. doi: 10.1016/j.revpalbo.2012.04.009.
- 27 508 Loisel, J. *et al.* (2014) ‘A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen  
28 509 accumulation’, *Holocene*, 24(9), pp. 1028–1042. doi: 10.1177/0959683614538073.
- 30 510 Loisel, J., Gallego-Sala, A. V. and Yu, Z. (2012) ‘Global-scale pattern of peatland Sphagnum growth driven by  
31 511 photosynthetically active radiation and growing season length’, *Biogeosciences*, 9(7), pp. 2737–2746. doi: 10.5194/bg-  
32 512 9-2737-2012.
- 33  
34 513 Loisel, J. and Garneau, M. (2010) ‘Late Holocene paleoecohydrology and carbon accumulation estimates from two  
35 514 boreal peat bogs in eastern Canada: Potential and limits of multi-proxy archives’, *Palaeogeography, Palaeoclimatology,*  
36 515 *Palaeoecology*. Elsevier B.V., 291(3–4), pp. 493–533. doi: 10.1016/j.palaeo.2010.03.020.
- 37 516 Loisel, J. and Yu, Z. (2013) ‘Surface vegetation patterning controls carbon accumulation in peatlands’, *Geophysical*  
38 517 *Research Letters*, 40(20), pp. 5508–5513. doi: 10.1002/grl.50744.
- 39  
40 518 Magnan, G. *et al.* (2018) ‘Impact of the Little Ice Age cooling and 20th century climate change on peatland vegetation  
41 519 dynamics in central and northern Alberta using a multi-proxy approach and high-resolution peat chronologies’,  
42 520 *Quaternary Science Reviews*. Elsevier Ltd, 185(March), pp. 230–243. doi: 10.1016/j.quascirev.2018.01.015.
- 43 521 Magnan, G. and Garneau, M. (2014a) ‘Climatic and autogenic control on Holocene carbon sequestration in  
44 522 ombrotrophic peatlands of maritime Quebec, eastern Canada’, *Holocene*, 24(9), pp. 1054–1062. doi:  
45 523 10.1177/0959683614540727.
- 46  
47 524 Magnan, G. and Garneau, M. (2014b) ‘Evaluating long-term regional climate variability in the maritime region of the St.  
48 525 Lawrence North Shore (eastern Canada) using a multi-site comparison of peat-based paleohydrological records’,  
49 526 *Journal of Quaternary Science*, 29(3), pp. 209–220. doi: 10.1002/jqs.2694.
- 50 527 Mathijssen, P. J. H. *et al.* (2016) ‘Reconstruction of Holocene carbon dynamics in a large boreal peatland complex,  
51 528 southern Finland’, *Quaternary Science Reviews*, 142, pp. 1–15. doi: 10.1016/j.quascirev.2016.04.013.
- 52  
53 529 Mathijssen, P. J. H. *et al.* (2017) ‘Lateral expansion and carbon exchange of a boreal peatland in Finland resulting in  
54 530 7000 years of positive radiative forcing’, *Journal of Geophysical Research: Biogeosciences*, 122(3), pp. 562–577. doi:  
55 531 10.1002/2016JG003749.
- 56  
57 532 Mauquoy, D. and van Geel, B. (2007) ‘Mire and Peat Macros’, (1935), pp. 537–554.
- 58 533 McGuire, A. D. *et al.* (2009) ‘Sensitivity of the carbon cycle in the Arctic to climate change’, *Ecological Monographs*,  
59 534 79(4), pp. 523–555. doi: 10.1890/08-2025.1.
- 60

- McGuire, A. D. *et al.* (2018) 'Dependence of the evolution of carbon dynamics in the northern permafrost region on the trajectory of climate change', *PNAS*, pp. 1–6. doi: 10.1073/pnas.1719903115.
- Naulier, M. *et al.* (2015) 'A millennial summer temperature reconstruction for northeastern Canada using oxygen isotopes in subfossil trees', *Climate of the Past*, 11(9), pp. 1153–1164. doi: 10.5194/cp-11-1153-2015.
- Nichols, J. E. *et al.* (2014) 'Impacts of climate and vegetation change on carbon accumulation in a south-central Alaskan peatland assessed with novel organic geochemical techniques', *The Holocene*, 24(9), pp. 1146–1155. doi: 10.1177/0959683614540729.
- Ovenden, L. (1990) 'Peat accumulation in northern wetlands', *Quaternary Research*, 33(3), pp. 377–386. doi: 10.1016/0033-5894(90)90063-Q.
- Packalen, M. S. and Finkelstein, S. A. (2014) 'Quantifying Holocene variability in carbon uptake and release since peat initiation in the Hudson Bay Lowlands, Canada', *Holocene*, 24(9), pp. 1063–1074. doi: 10.1177/0959683614540728.
- Packalen, M. S., Finkelstein, S. A. and McLaughlin, J. W. (2014) 'Carbon storage and potential methane production in the Hudson Bay Lowlands since mid-Holocene peat initiation', *Nature Communications*, 5(May), pp. 1–8. doi: 10.1038/ncomms5078.
- PAGES 2k Consortium (2013) 'Continental-scale temperature variability during the past two millennia', *Nature Geoscience*, 6(5), pp. 339–346. doi: 10.1038/ngeo1797.
- Payette, S. *et al.* (2004) 'Accelerated thawing of subarctic peatland permafrost over the last 50 years', *Geophysical Research Letters*, 31(18), pp. 1–4. doi: 10.1029/2004GL020358.
- van Der Plicht, J. (2004) 'Radiocarbon calibration - Past, present and future', *Nuclear Instruments and Methods in Physics Research, Section B: Beam Interactions with Materials and Atoms*, 223–224(SPEC. ISS.), pp. 353–358. doi: 10.1016/j.nimb.2004.04.069.
- Pratte, S., Garneau, M. and De Vleeschouwer, F. (2017) 'Increased atmospheric dust deposition during the Neoglacial in a boreal peat bog from north-eastern Canada', *Palaeogeography, Palaeoclimatology, Palaeoecology*, 469, pp. 34–46. doi: 10.1016/j.palaeo.2016.12.036.
- Reimer, P. (2013) 'IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP', *Radiocarbon*, 55(4), pp. 1869–1887. doi: 10.2458/azu\_js\_rc.55.16947.
- SOPFEU (Société de protection contre les feux) (2004) Mapping and Dating of Fires in Bay James Area, 1976–2004. Radisson/Val-d'Or office, Québec.
- Swindles, G. T. *et al.* (2012) 'Ecohydrological feedbacks confound peat-based climate reconstructions', *Geophysical Research Letters*, 39(11), pp. 2–5. doi: 10.1029/2012GL051500.
- Swindles, G. T. *et al.* (2015) 'The long-term fate of permafrost peatlands under rapid climate warming', *Scientific Reports*, 5. doi: 10.1038/srep17951.
- Thibault, S. and Payette, S. (2009) 'Recent Permafrost Degradation in Bogs of the James Bay Area, Northern Quebec, Canada', *Permafrost and Periglacial Processes*, 20(October), pp. 383–389. doi: 10.1002/ppp.
- Tivy, A. *et al.* (2011) 'Trends and variability in summer sea ice cover in the Canadian Arctic based on the Canadian Ice Service Digital Archive, 1960–2008 and 1968–2008', *Journal of Geophysical Research: Oceans*, 116(3). doi: 10.1029/2009JC005855.
- Treat, C. C. *et al.* (2016) 'Effects of permafrost aggradation on peat properties as determined from a pan-Arctic synthesis of plant macrofossils', *Journal of Geophysical Research: Biogeosciences*, 121(1), pp. 78–94. doi: 10.1002/2015JG003061.
- Tuittila, E.-S. *et al.* (2007) 'Quantifying patterns and controls of mire vegetation succession in a southern boreal bog in Finland using partial ordinations', *Journal of Vegetation Science*, 18(6), p. 891. doi: 10.1658/1100-9233(2007)18[891:QPACOM]2.0.CO;2.
- Tuittila, E. S. *et al.* (2012) 'Wetland chronosequence as a model of peatland development: Vegetation succession, peat and carbon accumulation', *Holocene*, 23(1), pp. 25–35. doi: 10.1177/0959683612450197.
- Turetsky, M., Manning, S. and Wieder R. (2004) 'Dating recent peat deposits', *Wetlands*, The Society of Wetland Scientists, 24(2): 324–356.

- Turunen, J. *et al.* (2002) 'Estimating carbon accumulation rates of undrained mires in Finland—application to boreal and subarctic regions', *The Holocene*, 12(1), pp. 69–80. doi: 10.1191/0959683602hl522rp.
- Turunen, J. (2003) 'Past and present carbon accumulation in undisturbed boreal and subarctic mires: A review' *Suo*, 54(1), pp. 15–28.
- Turunen, J. *et al.* (2004) 'Nitrogen deposition and increased carbon accumulation in ombrotrophic peatlands in eastern Canada', *Global Biogeochemical Cycles*, 18(3), pp. 1–12. doi: 10.1029/2003GB002154.
- Väliranta, M. *et al.* (2007) 'High-resolution reconstruction of wetness dynamics in a southern boreal raised bog, Finland, during the late Holocene: a quantitative approach', *Holocene*, 17(8), pp. 1093–1107. doi: 10.1177/0959683607082550.
- Väliranta, M. *et al.* (2017) 'Holocene fen–bog transitions, current status in Finland and future perspectives', *Holocene*, 27(5), pp. 752–764. doi: 10.1177/0959683616670471.
- Wilson, R. *et al.* (2016) 'Last millennium northern hemisphere summer temperatures from tree rings: Part I: The long term context', *Quaternary Science Reviews*, 134, pp. 1–18. doi: 10.1016/j.quascirev.2015.12.005.
- Yu, Z. *et al.* (2010) 'Global peatland dynamics since the Last Glacial Maximum', *Geophysical Research Letters*, 37(13), pp. 1–5. doi: 10.1029/2010GL043584.
- Yu, Z. (2011) 'Holocene carbon flux histories of the world's peatlands: Global carbon-cycle implications', *Holocene*, 21(5), pp. 761–774. doi: 10.1177/0959683610386982.
- Yu, Z., Beilman, D. W. and Jones, M. C. (2009) 'Sensitivity of Northern Peatland Carbon Dynamics to Holocene Climate Change', *Carbon Cycling in Northern Peatlands*, (C), pp. 55–69. doi: 10.1029/2008GM000822.
- Zhang, H. *et al.* (2018a) 'The role of climate change in regulating Arctic permafrost peatland hydrological and vegetation change over the last millennium', *Quaternary Science Reviews*, 182, pp. 121–130. doi: 10.1016/j.quascirev.2018.01.003.
- Zhang, H. *et al.* (2018b) 'Inconsistent response of Arctic permafrost peatland carbon accumulation to warm climate phases', *Global Biogeochemical Cycles*, doi: 10.1029/2018GB005980.